

Keeping of Galapagos lava lizard (*Microlophus albemarlensis* (Baur, 1890)) at BION Terrarium Center

Ivan Neizhko¹, Oleksii Marushchak², Yuliia Troshchinkova³

¹ – Senior Keeper, BION Terrarium Center, Kyiv, Ukraine

² – Head, Research and Development Department, BION Terrarium Center, Kyiv,
Ukraine

³ – Keeper, Laboratory of Desert Reptiles, BION Terrarium Center, Kyiv, Ukraine

Microlophus albemarlensis also known as Galápagos lava lizard or Albemarle lava lizard, is endemic to the Galápagos Islands, where it occurs on several islands in the western archipelago: Isabela, Fernandina, Santiago, Santa Cruz and Santa Fe, Seymour, Baltra, Daphne Major, Plaza Sur and Rábida. Of all *Microlophus* species, this one is the most widespread while other species inhabit single islands (Swash et al., 2005). Some authors however, consider populations on Santiago, Santa Cruz, and Santa Fe (and associated small islands (Fig. 1) to be distinct species (*M. jacobi*, *M. indefatigabilis* and *M. barringtonensis*, respectively).



Figure 1. *Microlophus albemarlensis* natural distribution range.



Figure 2. *Microlophus albemarlensis* male and female.

The species is commonly attributed to the genus *Microlophus* but has been historically placed in the genus *Tropidurus* (Benavides et al., 2009).

Males are easy to sex being around 5-6 months old. They usually have dark bodies with yellow-green markings, spinal crests of keeled scales, considerably bigger heads and well seen black pattern on the throat (Fig. 2). Females are reddish brown in color without markings. Females also have red or orange cheek patches when they reach maturity during breeding seasons (Fig. 2). Cheeks become brighter and more saturated during breeding season and gestation. Like most other lizards, both sexes exhibit changes in color depending on mood. Coloration also depends on where the lizards live. Those that live on dark lava are usually darker than those that live on light sandy land. Males are generally 2 to 3 times larger than females (Fig. 3). Adult males have total length ranging from 22 - 25 cm (8.66–9.84 in) while females range from 17 - 20 cm (6.69–7.87 in) (Jackson, 1985; Stebbins, et al., 1967).



Figure 3. *Microlophus albemarlensis* general physical appearance.

M. albemarlensis inhabits neotropical biogeographic region's lowland dry zones with loose soil and dry leaf litter where the lizards are able to bury at night. They also prefer rocky areas to hide underneath stones, rocks and inside crevices to escape from predators or too hot sun. Their habitat is characterized by such flora as cacti and vineplants, which provide shelter from the sun and attract insects (Stebbins, et al., 1967). In general, lizards appear when the sun rises (as early as 6am). Some lizards may start to forage immediately. The activity of the lizards increases by mid-morning when the ground starts to heat up rapidly. However, by mid-day, when the temperature rises beyond tolerable limits, they retreat from exposed positions to seek shelter under rocks, leaf litter or plants. These lizards do not communicate using any form of vocal communication. Most of their communication is through visual display because vision is their best-developed sense. "Push up" displays are used to ward off intruders as well as courtship communication. Change of skin color can communicate the mood of the lizard

from fear to aggression. Other actions are used indicate the intentions of individuals. For example, females are known to turn their back to males, raise their tail and shake it from side to side as a sign of rejection to approaching males. It is also believed that females use odors to dissuade male lizards during breeding season. (Stebbins, et al., 1967; Carpenter, 1977)

Breeding stock of 6 juveniles (later (3:3) was officially imported from the Netherlands in 2017.

Keeping and breeding requirements

Keeping. We keep adult Galapagos lava lizards in horizontal terrariums that should be long enough to provide temperature gradient. We use terrariums of 140*40*40 cm (55*16*16 in) for 1 pair (1:1). We use 2-4 cm (0.79-1.57 in) layer of coarse sand as a substrate. Decoration includes several ceramic shelters, flat stones, snags and sticks (Fig. 4, 5). We keep them in pairs since animals of the same sex demonstrate aggression towards each other.



Figure 4. Decorated terrarium for *M. albemarlensis* breeding group (1.2).

Lighting. Simple incandescent lamps (40-60 W) are used for day lighting (12-14 hours). We use them for making basking place. UV-light is necessary.

Temperature. Ambient temperature should not exceed +24 - +26 °C (75.2 - 78.8 F) at daytime and not be lower than +22 °C (71.6 F) at nighttime. At basking place temperature is about +35 °C (95.0 F). A flat stone is placed under the basking lamp.

Humidity. Humidity level is 50-60% with light spraying of warm corner every day.

Water. Water bowl is obligatory.

Diet. 80% of their diet consists of insects: crickets (*Gryllus assimilis*, *Gryllus bimaculatus*, *Gryllus locorojo*) and cockroaches (*Shelfordella tartara*). Other part of a diet is a finely chopped salad. In summer it is made of dandelions, nettle, hemp,

wormwood, alfalfa, coltsfoot, knotweed, clover, flowers of acacia, roses, hibiscus, mallow, linden, and echinacea. In other seasons the salad is made of lettuce, Chinese cabbage, dill, cilantro, parsley, sprouts, basil, carrots (finely grated), arugula, celery (root and leaves), spinach, broccoli, cauliflower and young bean leaves. It is necessary to add dried ingredients from summer salad set. Each time we give as many variable ingredients as possible. Food (both insects and herbs) is offered every other day. We offer “Repashy” calcium-vitamin supplements for every feeding. Dry food remnants are removed from the terrarium.



Figure 5. Young specimen of *M. albemarlensis*.

References

1. Andy Swash; Rob Still; Ian Lewington (2005). Birds, Mammals, and Reptiles of the Galápagos Islands: An Identification Guide. Yale University Press. pp. 120–121.
2. Benavides E.; Baum R.; Snell H. M.; Snell H. L.; Sites, Jr., J. W. (2009). "Island Biogeography of Galápagos Lava Lizards (Tropiduridae: Microlophus): Species Diversity and Colonization of the Archipelago." (.pdf) *Evolution*, 63 (6): 1606–1626.
3. Carpenter C. 1977. The aggressive displays of three species of South American iguanid lizards of the genus *Tropidurus*. *Herpetologica*, 33/3: 285.
4. Stebbins, R., J. Lowenstein, N. Cohen. 1967. A field study of the lava lizard (*Tropidurus albemarlensis*) in the Galapagos Islands. *Ecology*, 48/5: 839.
5. Jackson, M. H. 1985. Galapagos: A Natural History Guide. The University of Calgary Press.
6. Prieto, A., J. Leon, O. Lara. 1976. Reproduction in the tropical lizard, *Tropidurus hispidus*. *Herpetologica*, 32/3: 319.
7. https://animaldiversity.org/accounts/Microlophus_albemarlensis/
8. <https://www.inaturalist.org/taxa/39081-Microlophus-albemarlensis>